Young with three distinct streaks on the back, the central one being rather the broadest; the upper part of the sides with black net-work leaving a series of round olive spots.

Inhab. Cuba.

This species differs from all the other species of the genus we have in the British Museum in colour as well as in the peculiarity of the band of granular sides between the lateral and gular shields.

"Found under large stones in woods, and always on the ground looking for insects under the brushwood."—W. S. M.

ALLIGATOR.

9. *Alligator fissipes.* Caiman fissipes, *Spix. Braz.* t. 3. Champsa fissipes, *Wagler Icon.* t. 17.

Alligator Sclerops, *Br. Max. Abild. Braz.* t. 69. Crocodilus Sclerops, *Schinz. Nal. Rept.* t. 12. copied from *Pr. Max.*

Alligator Cynocephalus, *Dum. and Bibr. Rept.* iii. 87.

"Occurs sometimes 25 feet long."—W. S. M.

TESTUDO.

10. *Gopher Tortoise,* Bartram. *Testudo Polyphemus.*

Inhab. Cuba.

Lives in domestication.

AMPHISBÆNA.

11. *Amphisbæna punctata,* Bell Zool. Journ. iii.

Inhab. Cuba.

XIV.—*Characters of Five new Species of* Orchidaceous *Plants from Dominica.* By Prof. LINDLEY.

IN a small but interesting collection of dried plants of *Orchidaceæ,* collected in the island of Dominica, for the possession of which I am indebted to Prof. Henslow, there are the following new species.

1. STELIS *scabrida*; folio oblongo bidentato marginato basi cuneato caule longiore, racemo filiformi stricto folio longiore, bracteis ovatis amplexicaulibus acutissimis, petalis vertice scabridis labelloque lævi truncatis, antherâ pubescente.

2. PLEUROTHALLIS *aristata**; cæspitosa, folio ovali acuto

* Since this was in type, I have found this plant described and figured, under the same name which had been given it by me, in the 2nd volume of this work, p. 329, t. 15, by Sir W. Hooker. There are however differences enough between the two definitions to induce me to let my own remain.

Young with three distinct streaks on the back, the central one being rather the broadest; the upper part of the sides with black net-work leaving a series of round olive spots.

Inhab. Cuba.

This species differs from all the other species of the genus we have in the British Museum in colour as well as in the peculiarity of the band of granular sides between the lateral and gular shields.

"Found under large stones in woods, and always on the ground looking for insects under the brushwood."—W. S. M.

ALLIGATOR.

9. *Alligator fissipes.* Caiman fissipes, *Spix. Braz.* t. 3. Champsa fissipes, *Wagler Icon.* t. 17.

Alligator Sclerops, *Br. Max. Abild. Braz.* t. 69. Crocodilus Sclerops, *Schinz. Nal. Rept.* t. 12. copied from *Pr. Max.*

Alligator Cynocephalus, *Dum. and Bibr. Rept.* iii. 87.

"Occurs sometimes 25 feet long."—W. S. M.

TESTUDO.

10. *Gopher Tortoise,* Bartram. *Testudo Polyphemus.*

Inhab. Cuba.

Lives in domestication.

AMPHISBÆNA.

11. *Amphisbæna punctata,* Bell Zool. Journ. iii.

Inhab. Cuba.

XIV.—*Characters of Five new Species of* Orchidaceous *Plants from Dominica.* By Prof. LINDLEY.

IN a small but interesting collection of dried plants of *Orchidaceæ,* collected in the island of Dominica, for the possession of which I am indebted to Prof. Henslow, there are the following new species.

1. STELIS *scabrida*; folio oblongo bidentato marginato basi cuneato caule longiore, racemo filiformi stricto folio longiore, bracteis ovatis amplexicaulibus acutissimis, petalis vertice scabridis labelloque lævi truncatis, antherâ pubescente.

2. PLEUROTHALLIS *aristata**; cæspitosa, folio ovali acuto

* Since this was in type, I have found this plant described and figured, under the same name which had been given it by me, in the 2nd volume of this work, p. 329, t. 15, by Sir W. Hooker. There are however differences enough between the two definitions to induce me to let my own remain.

Young with three distinct streaks on the back, the central one being rather the broadest; the upper part of the sides with black net-work leaving a series of round olive spots.

Inhab. Cuba.

This species differs from all the other species of the genus we have in the British Museum in colour as well as in the peculiarity of the band of granular sides between the lateral and gular shields.

"Found under large stones in woods, and always on the ground looking for insects under the brushwood."—W. S. M.

ALLIGATOR.

9. *Alligator fissipes.* Caiman fissipes, *Spix. Braz.* t. 3. Champsa fissipes, *Wagler Icon.* t. 17.

Alligator Sclerops, *Br. Max. Abild. Braz.* t. 69. Crocodilus Sclerops, *Schinz. Nal. Rept.* t. 12. copied from *Pr. Max.*

Alligator Cynocephalus, *Dum. and Bibr. Rept.* iii. 87.

"Occurs sometimes 25 feet long."—W. S. M.

TESTUDO.

10. *Gopher Tortoise,* Bartram. *Testudo Polyphemus.*

Inhab. Cuba.

Lives in domestication.

AMPHISBÆNA.

11. *Amphisbæna punctata,* Bell Zool. Journ. iii.

Inhab. Cuba.

XIV.—*Characters of Five new Species of* Orchidaceous *Plants from Dominica.* By Prof. LINDLEY.

IN a small but interesting collection of dried plants of *Orchidaceæ,* collected in the island of Dominica, for the possession of which I am indebted to Prof. Henslow, there are the following new species.

1. STELIS *scabrida*; folio oblongo bidentato marginato basi cuneato caule longiore, racemo filiformi stricto folio longiore, bracteis ovatis amplexicaulibus acutissimis, petalis vertice scabridis labelloque lævi truncatis, antherâ pubescente.

2. PLEUROTHALLIS *aristata**; cæspitosa, folio ovali acuto

* Since this was in type, I have found this plant described and figured, under the same name which had been given it by me, in the 2nd volume of this work, p. 329, t. 15, by Sir W. Hooker. There are however differences enough between the two definitions to induce me to let my own remain.

apiculato basi angustato caule longiore, racemo capillari flexuoso erecto 4—6-floro folio ter longiore, bracteis vaginantibus cuspidatis, pedicellis elongatis, sepalis aristatis subciliatis lateralibus semiconnatis, petalis lanceolatis acuminatis semipinnatifidis duplò brevioribus, labello lineari glanduloso-hispido apice rotundato basi glabro utrinque auriculato.—Flowers purple, apparently streaked with yellow.

3. Camaridium *inflexum*; caule elongato subramoso pseudobulbifero, foliis geminis inæqualibus coriaceis oblongis basi canaliculatis, floribus solitariis, sepalis petalisque ovatis acutis, labello concavo erecto trilobo: laciniis membranaceis lateralibus inflexis falcatis serrulatis intermediâ subrotundâ leviter emarginatâ basi tuberculatâ.—Leaves about 2 inches long.

4. Rodriguezia *cochlearis*; pseudobulbis subrotundis compressis monophyllis cæspitosis, foliis oblongis sessilibus horizontalibus racemo erecto radicali subcorymboso paucifloro duplò brevioribus, sepalo supremo cum petalis obtuso lateralibus acutis semiconnatis breviore, labello obovato cochleari obtuso basi cristâ hippocrepicâ utrinque elongatâ bicarinato.—Flowers apparently yellow.

5. Pelexia *bursaria*; caule elongato folioso, foliis ovato-lanceolatis petiolatis, petiolis basi vaginantibus, scapo rachi ovariisque pilosis, bracteis membranaceis acuminatis pilosis ovarii longitudine, labello indiviso apice glanduloso, calcare inflato obtuso ovario paulò breviore.—A fine species with the habit of *Goodyera procera*. Stem from 1—1½ feet high.

XV.—*Information respecting Botanical and Zoological Travellers.*

Mr. Gould's Expedition to examine the Zoology of Australia.

We have received a letter from Mr. Gould written from the banks of the Hunter river; and having his permission to make such extracts from it as we may consider interesting to our readers, it gives us sincere pleasure to be able to state that the expedition of this orni-

apiculato basi angustato caule longiore, racemo capillari flexuoso erecto 4—6-floro folio ter longiore, bracteis vaginantibus cuspidatis, pedicellis elongatis, sepalis aristatis subciliatis lateralibus semiconnatis, petalis lanceolatis acuminatis semipinnatifidis duplò brevioribus, labello lineari glanduloso-hispido apice rotundato basi glabro utrinque auriculato.—Flowers purple, apparently streaked with yellow.

3. Camaridium *inflexum*; caule elongato subramoso pseudobulbifero, foliis geminis inæqualibus coriaceis oblongis basi canaliculatis, floribus solitariis, sepalis petalisque ovatis acutis, labello concavo erecto trilobo: laciniis membranaceis lateralibus inflexis falcatis serrulatis intermediâ subrotundâ leviter emarginatâ basi tuberculatâ.—Leaves about 2 inches long.

4. Rodriguezia *cochlearis*; pseudobulbis subrotundis compressis monophyllis cæspitosis, foliis oblongis sessilibus horizontalibus racemo erecto radicali subcorymboso paucifloro duplò brevioribus, sepalo supremo cum petalis obtuso lateralibus acutis semiconnatis breviore, labello obovato cochleari obtuso basi cristâ hippocrepicâ utrinque elongatâ bicarinato.—Flowers apparently yellow.

5. Pelexia *bursaria*; caule elongato folioso, foliis ovato-lanceolatis petiolatis, petiolis basi vaginantibus, scapo rachi ovariisque pilosis, bracteis membranaceis acuminatis pilosis ovarii longitudine, labello indiviso apice glanduloso, calcare inflato obtuso ovario paulò breviore.—A fine species with the habit of *Goodyera procera*. Stem from 1—1½ feet high.

XV.—*Information respecting Botanical and Zoological Travellers.*

Mr. Gould's Expedition to examine the Zoology of Australia.

We have received a letter from Mr. Gould written from the banks of the Hunter river; and having his permission to make such extracts from it as we may consider interesting to our readers, it gives us sincere pleasure to be able to state that the expedition of this orni-

apiculato basi angustato caule longiore, racemo capillari flexuoso erecto 4—6-floro folio ter longiore, bracteis vaginantibus cuspidatis, pedicellis elongatis, sepalis aristatis subciliatis lateralibus semiconnatis, petalis lanceolatis acuminatis semipinnatifidis duplò brevioribus, labello lineari glanduloso-hispido apice rotundato basi glabro utrinque auriculato.—Flowers purple, apparently streaked with yellow.

3. Camaridium *inflexum*; caule elongato subramoso pseudobulbifero, foliis geminis inæqualibus coriaceis oblongis basi canaliculatis, floribus solitariis, sepalis petalisque ovatis acutis, labello concavo erecto trilobo: laciniis membranaceis lateralibus inflexis falcatis serrulatis intermediâ subrotundâ leviter emarginatâ basi tuberculatâ.—Leaves about 2 inches long.

4. Rodriguezia *cochlearis*; pseudobulbis subrotundis compressis monophyllis cæspitosis, foliis oblongis sessilibus horizontalibus racemo erecto radicali subcorymboso paucifloro duplò brevioribus, sepalo supremo cum petalis obtuso lateralibus acutis semiconnatis breviore, labello obovato cochleari obtuso basi cristâ hippocrepicâ utrinque elongatâ bicarinato.—Flowers apparently yellow.

5. Pelexia *bursaria*; caule elongato folioso, foliis ovato-lanceolatis petiolatis, petiolis basi vaginantibus, scapo rachi ovariisque pilosis, bracteis membranaceis acuminatis pilosis ovarii longitudine, labello indiviso apice glanduloso, calcare inflato obtuso ovario paulò breviore.—A fine species with the habit of *Goodyera procera*. Stem from 1—1½ feet high.

XV.—*Information respecting Botanical and Zoological Travellers.*

Mr. Gould's Expedition to examine the Zoology of Australia.

We have received a letter from Mr. Gould written from the banks of the Hunter river; and having his permission to make such extracts from it as we may consider interesting to our readers, it gives us sincere pleasure to be able to state that the expedition of this orni-